arXiv:1212.2790v1 [math.CA] 12 Dec 2012

A Sturm-Liouville problem with a discontinuous coefficient and containing an eigenparameter in the boundary condition

Erdoğan Şen

Department of Mathematics, Faculty of Science and Letters, Namık Kemal University, 59030 Tekirdağ, Turkey

Department of Mathematics Engineering, Istanbul Technical University, Maslak, 34469 Istanbul, Turkey

e-mails: erdogan.math@gmail.com; koruc@itu.edu.tr

MSC (2010): 34B27, 34L10, 34L20.

Keywords : Sturm-Liouville problems; eigenparameter-dependent boundary condition; transmission conditions; asymptotics of eigenvalues and eigenfunctions;

ABSTRACT

In this paper, we study a Sturm-Liouville operator with eigenparameter-dependent boundary conditions and transmission conditions at three interior points. We give an operator-theoretical formulation, construct fundamental solutions and investigate some properties of the eigenvalues and corresponding eigenfunctions of the discontinuous Sturm-Liouville problem.

1 Introduction

In recent years, more and more researchers are interested in the discontinuous Sturm-Liouville problem for its application in physics (see [1-15]). Such problems are connected with discontinuous material properties, such as heat and mass transfer, varied assortment of physical transfer problems, vibrating string problems when the string loaded additionally with point masses, and diffraction problems [8,9]. Moreover there has been a growing interest in Sturm-Liouville problems with eigenparameter-dependent boundary conditions, i.e., the eigenparameter appears not only in the differential equations but also in the boundary conditions of the problems (see [1, 4-7, 10-13] and corresponding bibliography).

In this study, we examine eigenvalues and eigenfunctions of the differential equation

$$\tau u := -a(x)u''(x) + q(x)u(x) = \lambda u(x) \tag{1}$$

on $[a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$, with boundary conditions

$$\tau_1 u := \alpha_1 u\,(a) + \alpha_2 u'(a) = 0, \tag{2}$$

$$\tau_2 u := (\beta_1' \lambda + \beta_1)\, u\,(b) - (\beta_2' \lambda + \beta_2)\, u'(b) = 0 \tag{3}$$

and transmission conditions at the points of discontinuity $x = h_i$ $(i = 1, 2, 3)$,

$$\tau_3 u := \gamma_1\,(h_1 - 0) - \delta_1 u\,(h_1 + 0) = 0, \tag{4}$$

$$\tau_4 u := \gamma_1'\,(h_1 - 0) - \delta_1' u\,(h_1 + 0) = 0, \tag{5}$$

$$\tau_5 u := \gamma_2\,(h_2 - 0) - \delta_2 u\,(h_2 + 0) = 0, \tag{6}$$

$$\tau_6 u := \gamma_2'\,(h_2 - 0) - \delta_2' u\,(h_2 + 0) = 0, \tag{7}$$

$$\tau_7 u := \gamma_3\,(h_3 - 0) - \delta_3 u\,(h_3 + 0) = 0, \tag{8}$$

$$\tau_8 u := \gamma_3'\,(h_3 - 0) - \delta_3' u\,(h_3 + 0) = 0, \tag{9}$$

where $a(x) = a_1^2$, for $x \in [a, h_1)$, $a(x) = a_2^2$, for $x \in (h_1, h_2)$, $a(x) = a_3^2$ for $x \in (h_2, h_3)$, $a(x) = a_4^2$ for $x \in (h_3, b]$, $a < h_1 < h_2 < h_3 < b$, $q(x)$ is a given real-valued function continuous in $[a, h_1), (h_1, h_2), (h_2, h_3), (h_3, b]$ and has finite limits $q(h_i \pm 0) = \lim_{x \to h_i \pm 0} q(x)$ $(i = 1, 2, 3)$; λ is a complex eigenvalue parameter; $\gamma_i, \delta_i, \gamma_i', \delta_i'$ $(i = 1, 2, 3)$, $\alpha_j, \alpha_j', \beta_j, \beta_j'$ $(j = 1, 2)$ are real numbers; $|\alpha_1| + |\alpha_2| \neq 0$, $|\gamma_i| + |\delta_i| \neq 0$ and $|\gamma_i'| + |\delta_i'| \neq 0$ $(i = 1, 2, 3)$ and $\rho = \beta_1' \beta_2 - \beta_1 \beta_2' > 0$.

2 Operator formulation and some basic properties of this operator

In the Hilbert space $H = L_2\,(a, b) \oplus \mathbb{C}$ we define an inner product by

$$\langle F, G \rangle := \frac{1}{a_1^2} \int_a^{h_1} f\,(x)\, \overline{g\,(x)} dx + \frac{\delta_1 \delta_1'}{a_2^2 \gamma_1 \gamma_1'} \int_{h_1}^{h_2} f\,(x)\, \overline{g\,(x)} dx + \frac{\delta_1 \delta_1' \delta_2 \delta_2'}{a_3^2 \gamma_1 \gamma_1' \gamma_2 \gamma_2'} \int_{h_2}^{h_3} f\,(x)\, \overline{g\,(x)} dx +$$

$$+ \frac{1}{a_4^2} \left(\prod_{i=1}^{3} \frac{\delta_i \delta_i'}{\gamma_i \gamma_i'} \right) \int_{h_3}^{b} f\,(x)\, \overline{g\,(x)} dx + \frac{\prod\limits_{i=1}^{3} \frac{\delta_i \delta_i'}{\gamma_i \gamma_i'}}{a_4^2 \rho} f_1 \overline{g_1}$$

for

$$F := \begin{pmatrix} f\,(x) \\ f_1 \end{pmatrix}, \; G := \begin{pmatrix} g\,(x) \\ g_1 \end{pmatrix} \in H.$$

Following [7] for convenience we put

$$R_1\,(u) := \beta_1 u\,(1) - \beta_2 u'\,(1),$$

$$R_1'\,(u) := \beta_1' u\,(1) - \beta_2' u'\,(1).$$

For functionals $f(x)$, which defined on $[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$ and has finite limits $f(h_i \pm 0) := \lim_{x\to h_i \pm 0} f(x)$ $(i=1,2,3)$. By $f_i(x)$ $(i=1,2,3,4)$ we denote the functions

$$f_1(x) := \begin{cases} f(x), & x\in[-1,h_1), \\ \lim_{x\to h_1-0} f(x), & x=h_1, \end{cases}$$

$$f_2(x) := \begin{cases} \lim_{x\to h_1+0} f(x), & x=h_1, \\ f(x), & x\in(h_1,h_2), \\ \lim_{x\to h_2-0} f(x), & x=h_2, \end{cases}$$

$$, f_3(x) := \begin{cases} \lim_{x\to h_2+0} f(x), & x=h_2, \\ f(x), & x\in(h_2,h_3), \\ \lim_{x\to h_3-0} f(x), & x=h_3, \end{cases}$$

$$f_4(x) := \begin{cases} \lim_{x\to h_3+0} f(x), & x=h_3, \\ f(x), & x\in(h_3,b] \end{cases}$$

which are defined on $\Omega_1 := [a,h_1]$, $\Omega_2 := [h_1,h_2]$, $\Omega_3 := [h_2,h_3]$, $\Omega_4 := [h_3,b]$ respectively. We can rewrite the considered problem $(1)-(9)$ in the operator formulation as

$$AF = \lambda F$$

where

$$F := \begin{pmatrix} f(x) \\ -R_1'(f) \end{pmatrix} \in D(A)$$

and with

$$D(A) := \Big\{ F\in H \,\Big|\, f_i(x),\ f_i^{'}(x) \text{ are absolutely continuous in } \Omega_i\ \left(i=\overline{1,4}\right),\ \tau f \in L^2[a,b],\ \tau_1 f = \tau_2 f = ... = \tau_8 f = 0 \text{ and } f_1 = R_1^{'}(f) \Big\}$$

and

$$AF = \begin{pmatrix} \tau f \\ -R_1(f) \end{pmatrix}.$$

Consequently, the problem $(1)-(9)$ can be considered as the eigenvalue problem for the operator A.

Theorem 2.1. *The linear operator A is symmetric in H.*

Proof. Let F, $G\in D(A)$. By two partial integrations, we get

$$\begin{aligned}\langle AF,G\rangle &= \langle F,AG\rangle + \left(W(f,\overline{g};h_1-0) - W(f,\overline{g};a)\right) \\ &+ \frac{\delta_1\delta_1'}{\gamma_1\gamma_1'}\left(W(f,\overline{g};h_2-0) - W(f,\overline{g};h_1+0)\right) \\ &+ \frac{\delta_1\delta_1'\delta_2\delta_2'}{\gamma_1\gamma_1'\gamma_2\gamma_2'}\left(W(f,\overline{g};h_3-0) - W(f,\overline{g};h_2+0)\right) +\end{aligned}$$

$$+\left(\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}\right)\left(W\left(f,\overline{g};b\right)-W\left(f,\overline{g};h_3+0\right)\right)$$

$$+\frac{\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}}{\rho}\left(R_1'\left(f\right)R_1\left(\overline{g}\right)-R_1\left(f\right)R_1'\left(\overline{g}\right)\right) \qquad (10)$$

where

$$W\left(f,\overline{g};x\right)=f\left(x\right)\overline{g}'\left(x\right)-f'\left(x\right)\overline{g}\left(x\right)$$

denotes the Wronskians of the functions f and $\overline{g}$. Since f and $\overline{g}$ satisfy the boundary condition (2), it follows that

$$W\left(f,\overline{g};a\right)=0. \qquad (11)$$

From the transmission conditions (4)-(9) we get

$$\left\{\begin{array}{c} W\left(f,\overline{g};h_1-0\right)=\frac{\delta_1\delta_1'}{\gamma_1\gamma_1'}W\left(f,\overline{g};h_1+0\right),\\ W\left(f,\overline{g};h_2-0\right)=\frac{\delta_1\delta_1'\delta_2\delta_2'}{\gamma_1\gamma_1'\gamma_2\gamma_2'}W\left(f,\overline{g};h_2+0\right),\\ W\left(f,\overline{g};h_3-0\right)=\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}W\left(f,\overline{g};h_m+0\right). \end{array}\right. \qquad (12)$$

Furthermore,

$$\begin{aligned} & R_1'\left(f\right)R_1\left(\overline{g}\right)-R_1\left(f\right)R_1'\left(\overline{g}\right)\\ &=\left(\beta_1'f\left(b\right)-\beta_2'f'\left(b\right)\right)\left(\beta_1\overline{g}\left(b\right)-\beta_2\overline{g}'\left(b\right)\right)-\left(\beta_1f\left(b\right)-\beta_2f'\left(b\right)\right)\left(\beta_1'\overline{g}\left(b\right)-\beta_2'\overline{g}\left(b\right)\right)\\ &=\left(\beta_2\beta_1'-\beta_2'\beta_1\right)f'\left(b\right)\overline{g}\left(b\right)+\left(\beta_2'\beta_1-\beta_2\beta_1'\right)f\left(b\right)\overline{g}'\left(b\right)\\ &=\rho\left(f'\left(b\right)\overline{g}\left(b\right)-f\left(b\right)\overline{g}'\left(b\right)\right)=-\rho W\left(f,g;b\right). \end{aligned} \qquad (13)$$

Finally, substituting (11)-(13) in (10) then we get

$$\langle AF,G\rangle=\langle F,AG\rangle \quad \left(F,G\in D\left(A\right)\right) \qquad (14)$$

■

Corollary 2.1. *All eigenvalues of the problem (1)-(9) are real.*

We can now assume that all eigenfunctions are real-valued.

Corollary 2.2. *If λ_1 and λ_2 are two different eigenvalues of the problem (1)-(9), then the corresponding eigenfunctions u_1 and u_2 of this problem satisfy the following equality*:

$$\frac{1}{a_1^2}\int_a^{h_1}u_1\left(x\right)u_2\left(x\right)dx+\frac{\delta_1\delta_1'}{a_2^2\gamma_1\gamma_1'}\int_{h_1}^{h_2}u_1\left(x\right)u_2\left(x\right)dx+\frac{\delta_1\delta_1'\delta_2\delta_2'}{a_3^2\gamma_1\gamma_1'\gamma_2\gamma_2'}\int_{h_2}^{h_3}u_1\left(x\right)u_2\left(x\right)dx$$

$$+\frac{1}{a_4^2}\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}\int_{h_3}^{b}u_1\left(x\right)u_2\left(x\right)dx=-\frac{\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}}{a_4^2\rho}R_1'\left(u_1\right)R_1'\left(u_2\right).$$

In fact this formula means the orthogonality of eigenfunctions u_1 *and* u_2 *in the Hilbert space* H.

We need the following lemma, which can be proved by the same technique as in Theorem 1.5 in [15].

Lemma 2.1. *Let the real-valued function* $q(x)$ *be continuous in* $[a,b]$ *and* $f(\lambda), g(\lambda)$ *are given entire functions. Then for any* $\lambda \in \mathbb{C}$ *the equation*

$$-u'' + q(x)u = \lambda u, \quad x \in [a,b]$$

has a unique solution $u = u(x,\lambda)$ *satisfies the initial conditions*

$$u(a) = f(\lambda), u'(a) = g(\lambda) \ (\text{or } u(b) = f(\lambda), u'(b) = g(\lambda)).$$

For each fixed $x \in [a,b]$, $u(x,\lambda)$ *is an entire function of* λ.

We shall define two solutions

$$\phi_\lambda(x) = \begin{cases} \phi_{1\lambda}(x), \ x \in [a,h_1), \\ \phi_{2\lambda}(x), \ x \in (h_1,h_2), \\ \phi_{3\lambda}(x), \ x \in (h_2,h_3), \\ \phi_{4\lambda}(x), \ x \in (h_3,b], \end{cases} \text{ and } \chi_\lambda(x) = \begin{cases} \chi_{1\lambda}(x), \ x \in [a,h_1), \\ \chi_{2\lambda}(x), \ x \in (h_1,h_2), \\ \chi_{3\lambda}(x), \ x \in (h_2,h_3), \\ \chi_{4\lambda}(x), \ x \in (h_3,b], \end{cases}$$

of the Equation (1) as follows: Let $\phi_{1\lambda}(x) := \phi_1(x,\lambda)$ be the solution of Equation (1) on $[a,h_1]$, which satisfies the initial conditions

$$u(a) = \alpha_2, \ u'(a) = -\alpha_1. \tag{15}$$

By virtue of Lemma 2.1, after defining this solution, we may define the solution $\phi_2(x,\lambda) := \phi_{2\lambda}(x)$ of Equation (1) on $[h_1,h_2]$ by means of the solution $\phi_1(x,\lambda)$ by the initial conditions

$$u(h_1) = \frac{\gamma_1}{\delta_1}\phi_1(h_1,\lambda), \ u'(h_1) = \frac{\gamma_1'}{\delta_1'}\phi_1'(h_1,\lambda). \tag{16}$$

After defining this solution, we may define the solution $\phi_3(x,\lambda) := \phi_{3\lambda}(x)$ of Equation (1) on $[h_2,h_3]$ by means of the solution $\phi_2(x,\lambda)$ by the initial conditions

$$u(h_2) = \frac{\gamma_2}{\delta_2}\phi_2(h_2,\lambda), \ u'(h_2) = \frac{\gamma_2'}{\delta_2'}\phi_2'(h_2,\lambda). \tag{17}$$

Continuing in this manner, we may define the solution $\phi_4(x,\lambda) := \phi_{4\lambda}(x)$ of Equation (1) on $[h_3,b]$ by means of the solution $\phi_m(x,\lambda)$ by the initial conditions

$$u(h_3) = \frac{\gamma_3}{\delta_3}\phi_3(h_3,\lambda), \ u'(h_m) = \frac{\gamma_3'}{\delta_3'}\phi_3'(h_3,\lambda). \tag{18}$$

Therefore, $\phi(x,\lambda)$ satisfies the Equation (1) on $[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$, the boundary condition (3), and the transmission conditions (4)-(9).

Analogically, first we define the solution $\chi_{4\lambda}(x) := \chi_4(x,\lambda)$ on $[h_3,b]$ by the initial conditions

$$u(b) = \beta_2'\lambda + \beta_2, \ u'(b) = \beta_1'\lambda + \beta_1. \tag{19}$$

Again, after defining this solution, we may define the solution $\chi_{3\lambda}(x) := \chi_3(x,\lambda)$ of the Equation (1) on $[h_2, h_3]$ by the initial conditions

$$u(h_3) = \frac{\delta_3}{\gamma_3}\chi_4(h_3,\lambda)\,,\ \ u'(h_3) = \frac{\delta_3'}{\gamma_3'}\chi_4'(h_3,\lambda)\,. \tag{20}$$

Again, after defining this solution, we may define the solution $\chi_{2\lambda}(x) := \chi_2(x,\lambda)$ of the Equation (1) on $[h_1, h_2]$ by the initial conditions

$$u(h_2) = \frac{\delta_2}{\gamma_2}\chi_3(h_2,\lambda)\,,\ \ u'(h_2) = \frac{\delta_2'}{\gamma_2'}\chi_3'(h_2,\lambda)\,. \tag{21}$$

Continuing in this manner, we may define the solution $\chi_{1\lambda}(x) := \chi_1(x,\lambda)$ of the Equation (1) on $[a, h_1]$ by the initial conditions

$$u(h_1) = \frac{\delta_1}{\gamma_1}\chi_2(h_1,\lambda)\,,\ \ u'(h_1) = \frac{\delta_1'}{\gamma_1'}\chi_2'(h_1,\lambda)\,. \tag{22}$$

Therefore, $\chi(x,\lambda)$ satisfies the Equation (1) on $[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$, the boundary condition (3), and the transmission conditions (4)-(9).

It is obvious that the Wronskians

$$\omega_i(\lambda) := W_\lambda(\phi_i,\chi_i;x) := \phi_i(x,\lambda)\chi_i'(x,\lambda) - \phi_i'(x,\lambda)\chi_i(x,\lambda)\,,\quad x\in\Omega_i\ \ \left(i=\overline{1,4}\right)$$

are independent of $x\in\Omega_i$ and entire functions.

Lemma 2.2. *For each* $\lambda\in\mathbb{C}$, $\omega_1(\lambda) = \frac{\delta_1\delta_1'}{\gamma_1\gamma_1'}\omega_2(\lambda) = \frac{\delta_1\delta_1'\delta_2\delta_2'}{\gamma_1\gamma_1'\gamma_2\gamma_2'}\omega_3(\lambda) = \left(\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}\right)\omega_4(\lambda)\,.$

Proof. By the means of (16)-(18) and (20)-(22), the short calculation gives

$$\begin{aligned}W_\lambda(\phi_1,\chi_1;h_1) &= \frac{\delta_1\delta_1'}{\gamma_1\gamma_1'}W_\lambda(\phi_2,\chi_2;h_1) = \frac{\delta_1\delta_1'}{\gamma_1\gamma_1'}W_\lambda(\phi_3,\chi_3;h_2) = \frac{\delta_1\delta_1'\delta_2\delta_2'}{\gamma_1\gamma_1'\gamma_2\gamma_2'}W_\lambda(\phi_3,\chi_3;h_2)\\ &= \frac{\delta_1\delta_1'\delta_2\delta_2'}{\gamma_1\gamma_1'\gamma_2\gamma_2'}W_\lambda(\phi_4,\chi_4;h_3) = \left(\prod_{i=1}^{3}\frac{\delta_i\delta_i'}{\gamma_i\gamma_i'}\right)W_\lambda(\phi_4,\chi_4;h_3)\end{aligned}$$

so $\omega_1(\lambda) = \delta_1^2\omega_2(\lambda) = \delta_1^2\delta_2^2\omega_3(\lambda) = \delta_1^2\delta_2^2\delta_3^2\omega_4(\lambda)\,.$ ■

Now we may introduce the characteristic function

$$\omega(\lambda) := \omega_1(\lambda) = \delta_1^2\omega_2(\lambda) = \delta_1^2\delta_2^2\omega_3(\lambda) = \delta_1^2\delta_2^2\delta_3^2\omega_4(\lambda)\,.$$

Theorem 2.2. *The eigenvalues of the problem (1)-(9) are the zeros of the function* $\omega(\lambda)$.

Proof. Let $\omega(\lambda_0) = 0$. Then $W_{\lambda_0}(\phi_1,\chi_1;x) = 0$ and therefore the functions $\phi_{1\lambda_0}(x)$ and $\chi_{1\lambda_0}(x)$ are linearly dependent, i.e.

$$\chi_{1\lambda_0}(x) = k_1\phi_{1\lambda_0}(x)\,,\quad x\in[a,h_1]$$

for some $k_1 \neq 0$. From this, it follows that $\chi(x,\lambda_0)$ satisfies also the first boundary condition (2), so $\chi(x,\lambda_0)$ is an eigenfunction of the problem $(1)-(9)$ corresponding to this eigenvalue λ_0.

Now we let $u_0(x)$ be any eigenfunction corresponding to eigenvalue λ_0, but $\omega(\lambda_0) \neq 0$. Then the functions $\phi_1, \chi_1,\ \phi_2, \chi_2, \phi_3, \chi_3, \phi_4, \chi_4$ would be linearly independent on $[a,h_1], [h_1,h_2], [h_2,h_3]$ and $[h_3,b]$ respectively. Therefore $u_0(x)$ may be represented in the following form

$$u_0(x) = \begin{cases} c_1\phi_1(x,\lambda_0) + c_2\chi_1(x,\lambda_0), & x \in [a,h_1), \\ c_3\phi_2(x,\lambda_0) + c_4\chi_2(x,\lambda_0), & x \in (h_1,h_2), \\ c_5\phi_3(x,\lambda_0) + c_6\chi_3(x,\lambda_0), & x \in (h_2,h_3), \\ c_7\phi_4(x,\lambda_0) + c_8\chi_4(x,\lambda_0), & x \in (h_3,b]. \end{cases}$$

where at least one of the constants $c_1,\ c_2, ..., c_8$ is not zero. Considering the equations

$$\tau_\upsilon(u_0(x)) = 0, \quad \upsilon = \overline{1,8} \tag{23}$$

as the homogenous system of linear equations of the variables $c_1,\ c_2, ..., c_8$ and taking (16)-(18) and (20)-(22) into account, it follows that the determinant of this system is different from zero. Therefore, the system (23) has only the trivial solution $c_i = 0\ \left(i = \overline{1,8}\right)$. Thus we get a contradiction, which completes the proof. ■

Lemma 2.3. *If $\lambda = \lambda_0$ is an eigenvalue, then $\phi(x,\lambda_0)$ and $\chi(x,\lambda_0)$ are linearly dependent.*

Proof. Let $\lambda = \lambda_0$ be an eigenvalue. Then by virtue of Theorem 2.2

$$W(\phi_{i\lambda_0}, \chi_{i\lambda_0}; x) = \omega_i(\lambda_0) = 0$$

and hence

$$\chi_{i\lambda_0}(x) = k_i\phi_{i\lambda_0}(x) \qquad \left(i = \overline{1,4}\right) \tag{24}$$

for some $k_1 \neq 0,\ k_2 \neq 0, k_3 \neq 0, k_4 \neq 0$, We must show that $k_1 = k_2 = k_3 = k_4$. Suppose, if possible, that $k_3 \neq k_4$.

Taking into account the definitions of the solutions $\phi_i(x,\lambda)$ and $\chi_i(x,\lambda)$ from the equalities (24), we have

$$\begin{aligned} \tau_7(\chi_{\lambda_0}) &= \chi_{\lambda_0}(h_3-0) - \frac{\delta_3}{\gamma_3}\chi_{\lambda_0}(h_3+0) = \chi_{3\lambda_0}(h_3) - \frac{\delta_3}{\gamma_3}\chi_{4\lambda_0}(h_3) \\ &= k_3\phi_3(h_3) - \frac{\delta_3}{\gamma_3}k_4\phi_4(h_3) = k_3\frac{\delta_3}{\gamma_3}\phi_4(h_3) - \frac{\delta_3}{\gamma_3}k_4\phi_4(h_3) \\ &= \frac{\delta_3}{\gamma_3}(k_3-k_4)\phi_4(h_3) = 0. \end{aligned}$$

since $\tau_7(\chi_{\lambda_0}) = 0$ and $\frac{\delta_3}{\gamma_3}(k_3-k_4) \neq 0$, it follows that

$$\phi_{4\lambda_0}(h_3) = 0. \tag{25}$$

By the same procedure from $\tau_8(\chi_{\lambda_0}) = 0$ we can derive that

$$\phi'_{4\lambda_0}(h_3) = 0. \tag{26}$$

From the fact that $\phi_{4\lambda_0}(x)$ is a solution of the differential Equation (1) on $[h_3, b]$ and satisfies the initial conditions (25) and (26), it follows that $\phi_{4\lambda_0}(x) = 0$ identically on $[h_3, b]$ because of the well-known existence and uniqueness theorem for the initial value problems of the ordinary linear differential equations. Making use of (16)-(18) and (25)-(26), we may also derive that

$$\phi_{3\lambda_0}(h_m) = \phi'_{3\lambda_0}(h_3) = 0$$

Continuing in this matter, we may also find that

$$\begin{cases} \phi_{3\lambda_0}(h_2) = \phi'_{3\lambda_0}(h_2) = 0. \\ \phi_{2\lambda_0}(h_1) = \phi'_{2\lambda_0}(h_1) = 0 \\ \phi_{1\lambda_0}(h_1) = \phi'_{1\lambda_0}(h_1) = 0. \end{cases}$$

identically on $[h_2, h_3], [h_1, h_2], [a, h_1]$ respectively. Hence $\phi(x, \lambda_0) = 0$ identically on $[a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$. But this contradicts with (15). Hence $k_3 = k_4$. Analogically we can prove that $k_2 = k_3$ and $k_1 = k_2$. ■

Corollary 2.3. *If $\lambda = \lambda_0$ is an eigenvalue, then both $\phi(x, \lambda_0)$ and $\chi(x, \lambda_0)$ are eigenfunctions corresponding to this eigenvalue.*

Lemma 2.4. *All eigenvalues λ_n are simple zeros of $\omega(\lambda)$.*

Proof. Using the well-known Lagrange's formula, it can be shown that

$$(\lambda - \lambda_n)\left(\frac{1}{a_1^2}\int_a^{h_1} \phi_\lambda(x)\,\phi_{\lambda_n}(x)\,dx + \frac{1}{a_2^2}\int_{h_1}^{h_2} \phi_\lambda(x)\,\phi_{\lambda_n}(x)\,dx + \frac{1}{a_3^2}\int_{h_2}^{h_3} \phi_\lambda(x)\,\phi_{\lambda_n}(x)\,dx \right.$$

$$\left. + \frac{1}{a_4^2}\int_{h_3}^{b} \phi_\lambda(x)\,\phi_{\lambda_n}(x)\,dx\right) = W(\phi_\lambda, \phi_{\lambda_n}; b) \tag{27}$$

for any λ. Recall that

$$\chi_{\lambda_n}(x) = k_n \phi_{\lambda_n}(x), \quad x \in [a, h_1) \cup (h_1, h_2) \cup (h_2, h_3) \cup (h_3, b]$$

for some $k_n \neq 0$, $n = 1, 2, \ldots$. Using this equality for the right side of (27), we have

$$\begin{aligned} W(\phi_\lambda, \phi_{\lambda_n}; b) &= \frac{1}{k_n} W(\phi_\lambda, \chi_{\lambda_n}; b) = \frac{1}{k_n}\left(\lambda_n R_1'(\phi_\lambda) + R_1(\phi_\lambda)\right) \\ &= \frac{1}{k_n}\left[\omega(\lambda) - (\lambda - \lambda_n) R_1'(\phi_\lambda)\right] \\ &= (\lambda - \lambda_n)\frac{1}{k_n}\left[\frac{\omega(\lambda)}{\lambda - \lambda_n} - R_1'(\phi_\lambda)\right]. \end{aligned} \tag{28}$$

Substituting this formula in (27) and letting $\lambda \to \lambda_n$, we get

$$\frac{1}{a_1^2}\int_a^{h_1} (\phi_{\lambda_n}(x))^2\,dx + \frac{1}{a_2^2}\int_{h_1}^{h_2} (\phi_{\lambda_n}(x))^2\,dx + \frac{1}{a_3^2}\int_{h_2}^{h_3} (\phi_{\lambda_n}(x))^2\,dx + \frac{1}{a_4^2}\int_{h_3}^{b} (\phi_{\lambda_n}(x))^2\,dx$$

$$= \frac{1}{k_n}\left(\omega'(\lambda_n) - R_1'(\phi_{\lambda_n})\right).$$

Now putting

$$R_1'\left(\phi_{\lambda_n}\right)=\frac{1}{k_n}R_1'\left(\chi_{\lambda_n}\right)=\frac{\rho}{k_n}$$

in (28) we get $\omega'\left(\lambda_n\right)\neq 0$. ■

3 Asymptotic approximate formulas of $\omega\left(\lambda\right)$ for four distinct cases

We start by proving some lemmas.

Lemma 3.1. *Let $\phi\left(x,\lambda\right)$ be the solutions of Equation (1) defined in Section 2, and let $\lambda=s^2$. Then the following integral equations hold for $k=0,1$:*

$$\begin{aligned}\phi_{1\lambda}^{(k)}\left(x\right)&=\alpha_2\left(\cos\frac{s\left(x-a\right)}{a_1}\right)^{(k)}-\alpha_1\frac{a_1}{s}\left(\sin\frac{s\left(x-a\right)}{a_1}\right)^{(k)}\\&\quad+\frac{1}{a_1 s}\int\limits_a^x\left(\sin\frac{s\left(x-y\right)}{a_1}\right)^{(k)}q\left(y\right)\phi_{1\lambda}\left(y\right)dy,\end{aligned}\tag{29}$$

$$\begin{aligned}\phi_{2\lambda}^{(k)}\left(x\right)&=\tfrac{\gamma_1}{\delta_1}\phi_{1\lambda}\left(h_1\right)\left(\cos\tfrac{s(x-h_1)}{a_2}\right)^{(k)}+\tfrac{a_2}{s}\tfrac{\gamma_1'}{\delta_1'}\phi_{1\lambda}'\left(h_1\right)\left(\sin\tfrac{s(x-h_1)}{a_2}\right)^{(k)}\\&\quad+\tfrac{1}{a_2 s}\int\limits_{h_1}^x\left(\sin\tfrac{s(x-y)}{a_2}\right)^{(k)}q\left(y\right)\phi_{2\lambda}\left(y\right)dy,\end{aligned}\tag{30}$$

$$\begin{aligned}\phi_{3\lambda}^{(k)}\left(x\right)&=\tfrac{\gamma_2}{\delta_2}\phi_{2\lambda}\left(h_2\right)\left(\cos\tfrac{s(x-h_2)}{a_3}\right)^{(k)}+\tfrac{a_3}{s}\tfrac{\gamma_2'}{\delta_2'}\phi_{2\lambda}'\left(h_2\right)\left(\sin\tfrac{s(x-h_2)}{a_3}\right)^{(k)}\\&\quad+\tfrac{1}{a_3 s}\int\limits_{h_2}^x\left(\sin\tfrac{s(x-y)}{a_3}\right)^{(k)}q\left(y\right)\phi_{2\lambda}\left(y\right)dy,\end{aligned}\tag{31}$$

$$\begin{aligned}\phi_{4\lambda}^{(k)}\left(x\right)&=\tfrac{\gamma_3}{\delta_3}\phi_{3\lambda}\left(h_3\right)\left(\cos\tfrac{s(x-h_3)}{a_4}\right)^{(k)}+\tfrac{a_4}{s}\tfrac{\gamma_3'}{\delta_3'}\phi_{3\lambda}'\left(h_3\right)\left(\sin\tfrac{s(x-h_3)}{a_4}\right)^{(k)}\\&\quad+\tfrac{1}{a_4 s}\int\limits_{h_3}^x\left(\sin\tfrac{s(x-y)}{a_4}\right)^{(k)}q\left(y\right)\phi_{4\lambda}\left(y\right)dy,\end{aligned}\tag{32}$$

where $\left(\cdot\right)^{(k)}=\frac{d^k}{dx^k}\left(\cdot\right)$.

Proof. It is enough to substitute $s^2\phi_{1\lambda}\left(y\right)+\phi_{1\lambda}''\left(y\right), s^2\phi_{2\lambda}\left(y\right)+\phi_{2\lambda}''\left(y\right), s^2\phi_{3\lambda}\left(y\right)+\phi_{3\lambda}''\left(y\right), s^2\phi_{4\lambda}\left(y\right)+\phi_{4\lambda}''\left(y\right)$ instead of $q\left(y\right)\phi_{1\lambda}\left(y\right)$, $q\left(y\right)\phi_{2\lambda}\left(y\right)$, $q\left(y\right)\phi_{3\lambda}\left(y\right)$, $q\left(y\right)\phi_{4\lambda}\left(y\right)$ in the integral terms of the (29)-(32), respectively, and integrate by parts twice. ■

Lemma 3.2. *Let $\lambda=s^2$, $\operatorname{Im}s=t$. Then the functions $\phi_{i\lambda}\left(x\right)$ have the following asymptotic formulas for $\left|\lambda\right|\to\infty$, which hold uniformly for $x\in\Omega_i$ $\big($for $i=\overline{1,4}$ and $k=0,1.\big)$:*

If $\alpha_2 \neq 0$

$$\phi_{1\lambda}^{(k)}(x) = \alpha_2 \left(\cos \frac{s(x-a)}{a_1} \right)^{(k)} + O\left(|s|^{k-2} \exp \frac{|t|(x-a)}{a_1} \right), \tag{33}$$

$$\begin{aligned} \phi_{2\lambda}^{(k)}(x) = {} & \alpha_2 \frac{\gamma_1}{\delta_1} \left(\cos \frac{s(x-h_1)}{a_2} \right)^{(k)} \cos \frac{s(h_1-a)}{a_1} \\ & - \alpha_2 \frac{a_2 \gamma_1'}{a_1 \delta_1'} \left(\sin \frac{s(x-h_1)}{a_2} \right)^{(k)} \sin \frac{s(h_1-a)}{a_1} \\ & + O\left(|s|^{k-1} \exp \frac{|t| \left[a_1(x-h_1) + a_2(h_1-a) \right]}{a_1 a_2} \right), \end{aligned} \tag{34}$$

$$\begin{aligned} \phi_{3\lambda}^{(k)}(x) = {} & \alpha_2 \frac{\gamma_1 \gamma_2}{\delta_1 \delta_2} \left(\cos \frac{s(x-h_2)}{a_3} \right)^{(k)} \cos \frac{s(h_2-h_1)}{a_2} \cos \frac{s(h_1-a)}{a_1} \\ & - \alpha_2 \frac{\gamma_1' \gamma_2}{\delta_1' \delta_2} \left(\cos \frac{s(x-h_2)}{a_3} \right)^{(k)} \sin \frac{s(h_2-h_1)}{a_2} \sin \frac{s(h_1-a)}{a_1} \\ & - \alpha_2 \frac{\gamma_1 \gamma_2'}{\delta_1 \delta_2'} \left(\sin \frac{s(x-h_2)}{a_3} \right)^{(k)} \sin \frac{s(h_2-h_1)}{a_2} \cos \frac{s(h_1-a)}{a_1} \\ & - \alpha_2 \frac{\gamma_1' \gamma_2'}{\delta_1' \delta_2'} \left(\sin \frac{s(x-h_2)}{a_3} \right)^{(k)} \cos \frac{s(h_2-h_1)}{a_2} \sin \frac{s(h_1-a)}{a_1} \\ & + O\left(|s|^{k-1} \exp \frac{|t| \left[a_1 a_2 (x-h_2) + a_1 a_3 (h_2-h_1) + a_2 a_3 (h_1-a) \right]}{a_1 a_2 a_3} \right), \end{aligned} \tag{35}$$

$$\begin{aligned}\phi^{(k)}_{4\lambda}(x) = {} & \alpha_2 \frac{\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)} \cos\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1'\gamma_2\gamma_3 a_2}{\delta_1'\delta_2\delta_3 a_1}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)} \cos\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1\gamma_2'\gamma_3 a_3}{\delta_1\delta_2'\delta_3 a_2}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)} \sin\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1'\gamma_2'\gamma_3 a_3}{\delta_1'\delta_2'\delta_3 a_1}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)} \sin\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1\gamma_2\gamma_3' a_4}{\delta_1\delta_2\delta_3' a_3}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)} \sin\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1'\gamma_2\gamma_3' a_2 a_4}{\delta_1'\delta_2\delta_3' a_3 a_1}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)} \sin\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1\gamma_2'\gamma_3' a_4}{\delta_1\delta_2'\delta_3' a_2}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)} \cos\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 \frac{\gamma_1'\gamma_2'\gamma_3' a_4}{\delta_1'\delta_2'\delta_3' a_1}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)} \cos\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& + O\left(|s|^{k-1}\exp\frac{|t|\left[a_1a_2a_3(x-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)\right]}{a_1a_2a_3a_4}\right).\end{aligned} \tag{36}$$

If $\alpha_2 = 0$

$$\phi^{(k)}_{1\lambda}(x) = -\frac{a_1\alpha_1}{s}\left(\sin\frac{s(x-a)}{a_1}\right)^{(k)} + O\left(|s|^{k-2}\exp\frac{|t|(x-a)}{a_1}\right), \tag{37}$$

$$\begin{aligned}\phi^{(k)}_{2\lambda}(x) = {} & -\frac{a_1\alpha_1}{s}\frac{\gamma_1}{\delta_1}\left(\cos\frac{s(x-h_1)}{a_2}\right)^{(k)}\sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_2\frac{a_2\gamma_1'}{a_1\delta_1'}\left(\sin\frac{s(x-h_1)}{a_2}\right)^{(k)}\cos\frac{s(h_1-a)}{a_1} \\
& + O\left(|s|^{k-2}\exp\frac{|t|\left[a_1(x-h_1)+a_2(h_1-a)\right]}{a_1}\right),\end{aligned} \tag{38}$$

$$
\begin{aligned}
\phi_{3\lambda}^{(k)}(x) = & -\alpha_1 \frac{\gamma_1\gamma_2 a_1}{s\delta_1\delta_2}\left(\cos\frac{s(x-h_2)}{a_3}\right)^{(k)}\cos\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1'\gamma_2 a_2}{s\delta_1'\delta_2}\left(\cos\frac{s(x-h_2)}{a_3}\right)^{(k)}\sin\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1\gamma_2' a_1 a_3}{s\delta_1\delta_2' a_2}\left(\sin\frac{s(x-h_2)}{a_3}\right)^{(k)}\sin\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1'\gamma_2' a_3}{s\delta_1'\delta_2'}\left(\sin\frac{s(x-h_2)}{a_3}\right)^{(k)}\cos\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& +O\left(|s|^{k-2}\exp\frac{|t|\left[a_1a_2(x-h_2)+a_1a_3(h_2-h_1)+a_2a_3(h_1-a)\right]}{a_1a_2a_3}\right),
\end{aligned}
\tag{39}
$$

$$
\begin{aligned}
\phi_{4\lambda}^{(k)}(x) = & -\alpha_1 \frac{\gamma_1\gamma_2\gamma_3 a_1}{s\delta_1\delta_2\delta_3}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)}\cos\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1'\gamma_2\gamma_3 a_2}{s\delta_1'\delta_2\delta_3}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)}\cos\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& +\alpha_1 \frac{\gamma_1\gamma_2'\gamma_3 a_3 a_1}{s\delta_1\delta_2'\delta_3 a_2}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)}\sin\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1'\gamma_2'\gamma_3 a_3}{s\delta_1'\delta_2'\delta_3}\left(\cos\frac{s(x-h_3)}{a_4}\right)^{(k)}\sin\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& +\alpha_1 \frac{\gamma_1\gamma_2\gamma_3' a_4 a_1}{s\delta_1\delta_2\delta_3' a_3}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)}\sin\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& +\alpha_1 \frac{\gamma_1'\gamma_2\gamma_3' a_2 a_4}{s\delta_1'\delta_2\delta_3' a_3}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)}\sin\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& +\alpha_1 \frac{\gamma_1\gamma_2'\gamma_3' a_1 a_4}{s\delta_1\delta_2'\delta_3' a_2}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)}\cos\frac{s(h_3-h_2)}{a_3}\sin\frac{s(h_2-h_1)}{a_2}\sin\frac{s(h_1-a)}{a_1} \\
& -\alpha_1 \frac{\gamma_1'\gamma_2'\gamma_3' a_4}{s\delta_1'\delta_2'\delta_3'}\left(\sin\frac{s(x-h_3)}{a_4}\right)^{(k)}\cos\frac{s(h_3-h_2)}{a_3}\cos\frac{s(h_2-h_1)}{a_2}\cos\frac{s(h_1-a)}{a_1} \\
& +O\left(|s|^{k-2}\exp\frac{|t|\left[a_1a_2a_3(x-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)\right]}{a_1a_2a_3a_4}\right).
\end{aligned}
\tag{40}
$$

Proof. We may prove these formulas using the similar method as [15, Lemma 1.7 p. 9-10]. ■

Theorem 3.1. *Let* $\lambda = s^2$, $t = \operatorname{Im} s$. *Then the characteristic function* $\omega(\lambda)$ *has the following asymptotic formulas*:

Case 1: *If* $\beta_2' \neq 0$, $\alpha_2 \neq 0$, *then*

$$
\begin{aligned}
\omega_4(\lambda) = {} & \alpha_2 s^3 \beta_2' \frac{\gamma_1\gamma_2\gamma_3}{a_4\delta_1\delta_2\delta_3} \sin\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 s^3 \beta_2' \frac{a_2\gamma_1'\gamma_2\gamma_3}{a_1a_4\delta_1'\delta_2\delta_3} \sin\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 s^3 \beta_2' \frac{a_3\gamma_1\gamma_2'\gamma_3}{a_2a_4\delta_1\delta_2'\delta_3} \sin\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_2 s^3 \beta_2' \frac{a_3\gamma_1'\gamma_2'\gamma_3}{a_1a_4\delta_1'\delta_2'\delta_3} \sin\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& + \alpha_2 s^3 \beta_2' \frac{\gamma_1\gamma_2\gamma_3'}{a_3\delta_1\delta_2\delta_3'} \cos\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& + \alpha_2 s^3 \beta_2' \frac{a_2\gamma_1'\gamma_2\gamma_3'}{a_1a_3\delta_1'\delta_2\delta_3'} \cos\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& + \alpha_2 s^3 \beta_2' \frac{\gamma_1\gamma_2'\gamma_3'}{a_2\delta_1\delta_2'\delta_3'} \cos\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& + \alpha_2 s^3 \beta_2' \frac{\gamma_1'\gamma_2'\gamma_3'}{a_1\delta_1'\delta_2'\delta_3'} \cos\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& + O\left(|s|^2 \exp\frac{|t|\left[a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)\right]}{a_1a_2a_3a_4}\right).
\end{aligned}
\tag{41}
$$

Case 2: *If* $\beta_2' \neq 0$, $\alpha_2 = 0$, *then*

$$
\begin{aligned}
\omega_4(\lambda) = {} & -\alpha_1 s^2 \beta_2' \frac{a_1\gamma_1\gamma_2\gamma_3}{a_4\delta_1\delta_2\delta_3} \sin\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_1 s^2 \beta_2' \frac{a_2\gamma_1'\gamma_2\gamma_3}{a_4\delta_1'\delta_2\delta_3} \sin\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_1 s^2 \beta_2' \frac{a_1a_3\gamma_1\gamma_2'\gamma_3}{a_2a_4\delta_1\delta_2'\delta_3} \sin\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& - \alpha_1 s^2 \beta_2' \frac{a_3\gamma_1'\gamma_2'\gamma_3}{a_4\delta_1'\delta_2'\delta_3} \sin\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_1 s^2 \beta_2' \frac{a_1\gamma_1\gamma_2\gamma_3'}{a_3\delta_1\delta_2\delta_3'} \cos\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& + \alpha_1 s^2 \beta_2' \frac{a_2\gamma_1'\gamma_2\gamma_3'}{a_3\delta_1'\delta_2\delta_3'} \cos\frac{s(b-h_3)}{a_4} \sin\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1} \\
& - \alpha_1 s^2 \beta_2' \frac{a_1\gamma_1\gamma_2'\gamma_3'}{a_2\delta_1\delta_2'\delta_3'} \cos\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \sin\frac{s(h_2-h_1)}{a_2} \sin\frac{s(h_1-a)}{a_1} \\
& + \alpha_1 s^2 \beta_2' \frac{\gamma_1'\gamma_2'\gamma_3'}{\delta_1'\delta_2'\delta_3'} \cos\frac{s(b-h_3)}{a_4} \cos\frac{s(h_3-h_2)}{a_3} \cos\frac{s(h_2-h_1)}{a_2} \cos\frac{s(h_1-a)}{a_1}
\end{aligned}
$$

$$+O\left(|s|\exp\frac{|t|\left[a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)\right]}{a_1a_2a_3a_4}\right). \tag{42}$$

Case 3: *If* $\beta_2' = 0,\ \alpha_2 \neq 0,$ *then*

$$\begin{aligned}
\omega_4\left(\lambda\right) = {} & \alpha_2 s^2\beta_1'\frac{\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_2 s^2\beta_1'\frac{a_2\gamma_1'\gamma_2\gamma_3}{a_1\delta_1'\delta_2\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_2 s^2\beta_1'\frac{a_3\gamma_1\gamma_2'\gamma_3}{a_2\delta_1\delta_2'\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_2 s^2\beta_1'\frac{a_3\gamma_1'\gamma_2'\gamma_3}{a_1\delta_1'\delta_2'\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_2 s^2\beta_1'\frac{a_4\gamma_1\gamma_2\gamma_3'}{a_3\delta_1\delta_2\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& +\alpha_2 s^2\beta_1'\frac{a_2a_4\gamma_1'\gamma_2\gamma_3'}{a_3a_1\delta_1'\delta_2\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_2 s^2\beta_1'\frac{a_4\gamma_1\gamma_2'\gamma_3'}{a_2\delta_1\delta_2'\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& +\alpha_2 s^2\beta_1'\frac{a_4\gamma_1'\gamma_2'\gamma_3'}{a_1\delta_1'\delta_2'\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}
\end{aligned}$$

$$+O\left(|s|\exp\frac{|t|\left[a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)\right]}{a_1a_2a_3a_4}\right). \tag{43}$$

Case 4: *If* $\beta_2' = 0,\ \alpha_2 = 0,$ *then*

$$\begin{aligned}
\omega_4\left(\lambda\right) = {} & -\alpha_1 s\beta_1'\frac{a_1\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_1 s\beta_1'\frac{a_2\gamma_1'\gamma_2\gamma_3}{\delta_1'\delta_2\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_1 s\beta_1'\frac{a_1a_3\gamma_1\gamma_2'\gamma_3}{a_2\delta_1\delta_2'\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_1 s\beta_1'\frac{a_3\gamma_1'\gamma_2'\gamma_3}{\delta_1'\delta_2'\delta_3}\cos\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_1 s\beta_1'\frac{a_1a_4\gamma_1\gamma_2\gamma_3'}{a_3\delta_1\delta_2\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& +\alpha_1 s\beta_1'\frac{a_2a_4\gamma_1'\gamma_2\gamma_3'}{a_3\delta_1'\delta_2\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\sin\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}\\
& -\alpha_1 s\beta_1'\frac{a_1a_4\gamma_1\gamma_2'\gamma_3'}{a_2\delta_1\delta_2'\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\sin\frac{s\left(h_2-h_1\right)}{a_2}\sin\frac{s\left(h_1-a\right)}{a_1}\\
& +\alpha_1 s\beta_1'\frac{a_4\gamma_1'\gamma_2'\gamma_3'}{\delta_1'\delta_2'\delta_3'}\sin\frac{s\left(b-h_3\right)}{a_4}\cos\frac{s\left(h_3-h_2\right)}{a_3}\cos\frac{s\left(h_2-h_1\right)}{a_2}\cos\frac{s\left(h_1-a\right)}{a_1}
\end{aligned}$$

$$+O\left(\exp\frac{|t|\left[a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)\right]}{a_1a_2a_3a_4}\right). \tag{44}$$

Proof. The proof is completed by substituting (36) and (40) into the representation

$$\begin{aligned}\omega_4(\lambda)&=\lambda\left[\beta_1'\phi_{4\lambda}(b)-\beta_2'\phi_{4\lambda}'(b)\right]+\left[\beta_1\phi_{4\lambda}(b)-\beta_2\phi_{4\lambda}'(b)\right]\\&=-\lambda\beta_2'\phi_{4\lambda}'(b)+\lambda\beta_1'\phi_{4\lambda}(b)+\beta_1\phi_{4\lambda}(b)-\beta_2'\phi_{4\lambda}'(b).\end{aligned} \tag{45}$$

■

Corollary 3.1. *The eigenvalues of the problem (1)-(9) are bounded below.*
Proof. Putting $s=it$ $(t>0)$ in the above formulas, it follows that $\omega_4\left(-t^2\right)\to\infty$ as $t\to\infty$. Therefore, $\omega_4(\lambda)\neq0$ for λ negative and sufficiently large. ■

4 Asymptotic formulas for eigenvalues and eigenfunctions

Now we can obtain the asymptotic approximation formulas for the eigenvalues of the considered problem (1)-(9).

Since the eigenvalues coincide with the zeros of the entire function $\omega_4(\lambda)$, it follows that they have no finite limit. Moreover, we know from Corollaries 2.1 and 3.1 that all eigenvalues are real and bounded below. Hence, we may renumber them as $\lambda_0\leq\lambda_1\leq\lambda_2\leq\ldots$, listed according to their multiplicity.

Theorem 4.1. *The eigenvalues* $\lambda_n=s_n^2$, $n=0,1,2,\ldots$ *of the problem (1)-(9) have the following asymptotic formulas for* $n\to\infty$:

Case 1: *If* $\beta_2^{'}\neq0,\alpha_2\neq0$, *then*

$$s_n=\frac{a_1a_2a_3a_4\pi\left(n-1\right)}{a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)}+O\left(\frac{1}{n}\right). \tag{46}$$

Case 2: *If* $\beta_2^{'}\neq0,\alpha_2=0$, *then*

$$s_n=\frac{a_1a_2a_3a_4\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)}+O\left(\frac{1}{n}\right). \tag{47}$$

Case 3: *If* $\beta_2^{'}=0,\alpha_2\neq0$, *then*

$$s_n=\frac{a_1a_2a_3a_4\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3\left(b-h_3\right)+a_1a_2a_4\left(h_3-h_2\right)+a_1a_3a_4\left(h_2-h_1\right)+a_2a_3a_4\left(h_1-a\right)}+O\left(\frac{1}{n}\right). \tag{48}$$

Case 4: *If* $\beta_2^{'} = 0, \alpha_2 = 0,$ *then*

$$s_n = \frac{a_1 a_2 a_3 a_4 \pi n}{a_1 a_2 a_3 (b - h_3) + a_1 a_2 a_4 (h_3 - h_2) + a_1 a_3 a_4 (h_2 - h_1) + a_2 a_3 a_4 (h_1 - a)} + O\left(\frac{1}{n}\right). \tag{49}$$

Proof. We shall only consider the first case. The other cases may be considered similarly. Denoting $\omega_0 (s)$ and $\omega_2 (s)$ the first and O-term of the right of (42) repectively, we shall apply the well-known Rouché's theorem, which asserts that if $f(s)$ and $g(s)$ are analytic inside and on a closed contour C, and $|g(s)| < |f(s)|$ on C, then $f(s)$ and $f(s) + g(s)$ have the same number zeros inside C, provided that each zero is counted according to their multiplicity. It is readily shown that $|\overline{\omega}_1 (s)| > |\overline{\omega}_2 (s)|$ on the contours

$$C_n := \left\{ s \in \mathbb{C} \, | |s| = \frac{\pi a_1 a_2 a_3 a_4 \left(n + \frac{1}{2}\right)}{a_1 a_2 a_3 (b - h_3) + a_1 a_2 a_4 (h_3 - h_2) + a_1 a_3 a_4 (h_2 - h_1) + a_2 a_3 a_4 (h_1 - a)} \right\}$$

for sufficiently large n.

Let $\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq ...$ be zeros of $\omega(\lambda)$ and $\lambda_n = s_n^2$. Since inside the contour C_n, $\overline{\omega}_1 (s)$ has zeros at points $s = 0$ with multiplicity 4 and

$$s = \frac{k \pi a_1 a_2 a_3 a_4}{2 \left[a_1 a_2 a_3 (b - h_3) + a_1 a_2 a_4 (h_3 - h_2) + a_1 a_3 a_4 (h_2 - h_1) + a_2 a_3 a_4 (h_1 - a)\right]},$$

$k = \pm 1, \pm 2, ..., \pm n$

with multiplicity 1, and the number of zeros is $2n + 4$, it follows that

$$s_n = \frac{\pi a_1 a_2 a_3 a_4 (n - 1)}{a_1 a_2 a_3 (b - h_3) + a_1 a_2 a_4 (h_3 - h_2) + a_1 a_3 a_4 (h_2 - h_1) + a_2 a_3 a_4 (h_1 - a)} + \delta_n,$$

where $\delta_n = O(1)$, more precisely

$$|\delta_n| < \frac{\pi a_1 a_2 a_3 a_4}{4 \left[a_1 a_2 a_3 (b - h_3) + a_1 a_2 a_4 (h_3 - h_2) + a_1 a_3 a_4 (h_2 - h_1) + a_2 a_3 a_4 (h_1 - a)\right]}$$

for sufficiently large n. By substituting this in (41), we derive that $\delta_n = O\left(\frac{1}{n}\right)$, which completes the proof. ■

Recalling that $\phi(x, \lambda_n)$ an eigenfunction according to eigenvalue λ_n, by putting (46)-(49) in the (33)-(40) for $k = 0$, we derive the eigenfunctions:

In case 1

$$\phi(x, \lambda_n) = \begin{cases} \alpha_2 \cos\left[\frac{a_2 a_3 a_4 (x-a) \pi (n-1)}{a_1 a_2 a_3 (b-h_3) + a_1 a_2 a_4 (h_3-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)}\right] \\ \quad + O\left(\frac{1}{n}\right), \quad x \in [a, h_1) \\ \frac{\alpha_2 \gamma_1}{\delta_1} \cos\left[\frac{[a_1 a_3 a_4 (x-h_1) + a_2 a_3 a_4 (h_1-a)] \pi (n-1)}{a_1 a_2 a_3 (b-h_3) + a_1 a_2 a_4 (h_3-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)}\right] \\ \quad + O\left(\frac{1}{n}\right), \quad x \in (h_1, h_2) \\ \frac{\alpha_2 \gamma_1 \gamma_2}{\delta_1 \delta_2} \cos\left[\frac{[a_1 a_2 a_3 (x-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)] \pi (n-1)}{a_1 a_2 a_3 (b-h_3) + a_1 a_2 a_4 (h_3-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)}\right] \\ \quad + O\left(\frac{1}{n}\right), \quad x \in (h_2, h_3) \\ \frac{\alpha_2 \gamma_1 \gamma_2 \gamma_3}{\delta_1 \delta_2 \delta_3} \cos\left[\frac{[a_1 a_2 a_4 (x-h_3) + a_1 a_2 a_3 (h_3-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)] \pi (n-1)}{a_1 a_2 a_3 (b-h_3) + a_1 a_2 a_4 (h_3-h_2) + a_1 a_3 a_4 (h_2-h_1) + a_2 a_3 a_4 (h_1-a)}\right] \\ \quad + O\left(\frac{1}{n}\right), \quad x \in (h_3, b] \end{cases}$$

In case 2

$$\phi\left(x,\lambda_n\right)=\begin{cases} -\alpha_1\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi\left(n-\frac{1}{2}\right)} \\ \times\sin\left[\frac{a_2a_3a_4(x-a)\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right), \\ x\in[a,h_1) \\ -\frac{\alpha_1\gamma_1}{\delta_1}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi\left(n-\frac{1}{2}\right)} \\ \times\sin\left[\frac{[a_1a_3a_4(x-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right), \\ x\in(h_1,h_2) \\ -\frac{\alpha_1\gamma_1\gamma_2}{\delta_1\delta_2}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi\left(n-\frac{1}{2}\right)} \\ \times\sin\left[\frac{[a_1a_2a_3(x-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right), \\ x\in(h_2,h_3) \\ -\frac{\alpha_1\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi\left(n-\frac{1}{2}\right)} \\ \times\sin\left[\frac{[a_1a_2a_4(x-h_3)+a_1a_2a_3(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right), \\ x\in(h_3,b] \end{cases}$$

In case 3

$$\phi\left(x,\lambda_n\right)=\begin{cases} \alpha_2\cos\left[\frac{a_2a_3a_4(x-a)\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right] \\ +O\left(\frac{1}{n}\right),\quad x\in[a,h_1) \\ \frac{\alpha_2\gamma_1}{\delta_1}\cos\left[\frac{[a_1a_3a_4(x-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right] \\ +O\left(\frac{1}{n}\right),\quad x\in(h_1,h_2) \\ \frac{\alpha_2\gamma_1\gamma_2}{\delta_1\delta_2}\cos\left[\frac{[a_1a_2a_3(x-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right] \\ +O\left(\frac{1}{n}\right),\quad x\in(h_2,h_3) \\ \frac{\alpha_2\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\cos\left[\frac{[a_1a_2a_4(x-h_3)+a_1a_2a_3(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi\left(n-\frac{1}{2}\right)}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right] \\ +O\left(\frac{1}{n}\right),\quad x\in(h_3,b] \end{cases}$$

In case 4

$$\phi\left(x,\lambda_n\right)=\begin{cases}\begin{array}{c}-\alpha_1\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi n}\\ \times\sin\left[\frac{a_2a_3a_4(x-a)\pi n}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right),\\ x\in[a,h_1)\end{array}\\ \begin{array}{c}-\frac{\alpha_1\gamma_1}{\delta_1}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi n}\\ \times\sin\left[\frac{[a_1a_3a_4(x-h_1)+a_2a_3a_4(h_1-a)]\pi n}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right),\\ x\in(h_1,h_2)\end{array}\\ \begin{array}{c}-\frac{\alpha_1\gamma_1\gamma_2}{\delta_1\delta_2}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi n}\\ \times\sin\left[\frac{[a_1a_2a_3(x-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi n}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right),\\ x\in(h_2,h_3)\end{array}\\ \begin{array}{c}-\frac{\alpha_1\gamma_1\gamma_2\gamma_3}{\delta_1\delta_2\delta_3}\frac{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}{a_2a_3a_4\pi n}\\ \times\sin\left[\frac{[a_1a_2a_4(x-h_3)+a_1a_2a_3(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)]\pi n}{a_1a_2a_3(b-h_3)+a_1a_2a_4(h_3-h_2)+a_1a_3a_4(h_2-h_1)+a_2a_3a_4(h_1-a)}\right]+O\left(\frac{1}{n^2}\right),\\ x\in(h_3,b]\end{array}\end{cases}$$

All these asymptotic formulas hold uniformly for $x\in[a,h_1)\cup(h_1,h_2)\cup(h_2,h_3)\cup(h_3,b]$.

References

[1] J. Walter, Regular eigenvalue problems with eigenvalue parameter in the boundary conditions, Math. Z. 133 (1973) 301–312.

[2] S. Yakubov, Y. Yakubov, Abel basis of root functions of regular boundary value problems, Math. Nachr. 197 (1999) 157–187.

[3] S. Yakubov, Y. Yakubov, Differential-Operator Equations. Ordinary and Partial Differential Equations, Chapman and Hall/CRC.

[4] N. B. Kerimov, Kh. R. Mamedov, On a boundary value problem with a spectral parameter in the boundary conditions, Sibirsk. Mat. Zh. 40 (2) (1999) 325–335 (English translation: Siberian Math. J. 40 (2) (1999) 281–290).

[5] M. Kadakal and O. Sh. Mukhtarov, Eigenvalues and eigenfunctions of Sturm-Liouville problems with discontinuities at two points, Scientia Magna 6 (4) (2010) 33-48.

[6] Q. Yang and W. Wang, Asymptotic behavior of a differential operator with two point discontinuities containing eigenparameter-dependent boundary conditions, Mathematical Methods in the applied sciences 34 (2011) 373-383.

[7] C. T. Fulton, Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions, Proc. Roy. Soc. Edinburgh Sect. A 77 (1977) 293–308.

[8] M. Kobayashi, Eigenvalues of discontinuous Sturm–Liouville problems with symmetric potential, Comput. Math. Appl. 18 (4) (1989) 357–364.

[9] A. V. Likov, Yu. A. Mikhailov, The Theory of Heat and Mass Transfer, Qosenerqoizdat (1963) (in Russian).

[10] P. A. Binding, P. J. Browne, B. A. Watson, Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter II. J. Comput. Appl. Math. 148 (2002) 147-169.

[11] P. A. Binding, R. Hryniv, H. Langer, Elliptic eigenvalue problems with eigenparameter dependent boundary conditions, J. Diff. Eq., 174 (2001) 30-54.

[12] A. A. Shkalikov, Boundary value problems for ordinary differential equations with a parameter in boundary condition, Trudy Sem. Imeny I. G. Petrowsgo 9 (1983) 190–229.

[13] E. Şen and A. Bayramov, Calculation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument which contains a spectral parameter in the boundary condition, Mathematical and Computer Modelling, 54 (2011) 3090-3097.

[14] E. Şen and A. Bayramov, Asymptotic formulations of the eigenvalues and eigenfunctions for a boundary value problem, Mathematical Methods in the Applied Sciences, doi: 10.1002/mma.2699 (2012).

[15] E. C. Titchmarsh, Eigenfunctions Expansion Associated With Second Order Differential Equation 1, 2nd ed., Oxford University Press, London, 1962.